

## INTRODUCTION

Coronary artery disease (CAD) is a disease characterized by narrowing of the coronary arteries due to accumulation of plaque, cholesterol, and calcium within the walls of the coronary arteries [1]. Cardiovascular disease was the number one cause of deaths in 2014 accounting for 30% of all death cases in the UAE [2]. While invasive imaging procedures remain unreplacable in diagnosing and treating stenosis in CAD patients, a need arises for a noninvasive imaging technique for preventive purposes in asymptomatic or minimally symptomatic patients [3].

## OBJECTIVE

The primary aim of this study is to evaluate stenosis using Coronary Computed Tomography Angiography (CCTA) and Computed Tomography Calcium Scoring (CTCS), and correlate the findings with Invasive Coronary Angiography (ICA) in symptomatic Coronary Artery Disease (CAD) patients.

## MATERIALS & METHODS

The study included 24 patients (14 males and 10 females: mean age 62.3 years ranging from 34–74, with known CAD. The data was retrieved from the Radiology Department of University Hospital in Sharjah and was retrospectively reviewed. All patients have undergone ICA, 16 undergone CCTA, and 22 undergone CTCS. CCTA and CTCS were performed using Multi Detector Computed Tomography (MDCT), Fig 2. CCTA and ICA stenosis measurement was done using the Quantitative Coronary Angiography (QCA) method, Fig 1.




Fig 1, QCA stenosis measurement in CCTA showing severity and type of plaque present in a defined ROI.




Fig 2, An axial view of the LAD showing multiple calcification plaques in CTCS (A) and CCTA (B) and calcium score results (C) using MDCT.

### Statistical Analysis

All the statistical analysis was performed using GraphPad Prism 7 (version 7.03) as well as IBM SPSS (version 22) statistical software.

## RESULTS & DISCUSSION

### Finding 1: CCTA vs ICA

Table 1: Segment-based ( $\geq 50\%$ ) diagnostic accuracy of MDCT showing an underestimation of significant stenosis in 13%, and an overestimation in 10% of patients comparing to ICA.

|                |       |
|----------------|-------|
| True positive  | 13    |
| True negative  | 136   |
| False positive | 20    |
| False negative | 26    |
| Sensitivity    | 39.4% |
| specificity    | 84.0% |

Correlation of ICA & CCTA Segment-based  
( $r = 0.2595$ )




Fig 3: Spearman rank correlation graph showing significantly ( $P=0.0002$ ) weak correlation between CCTA and ICA.




Fig 4: CCTA image showing calcific plaques at the mid portion of LCX and origin of OM (A), ICA image shows narrowing of the mid portion of LCX which indicates stenosis (B).

Table 2, Vessel-based diagnostic accuracy of MDCT showing LAD to have the lowest detection specificity and LM to have the highest.

|     | TP | TN | FP | FN | Sensitivity | Specificity |
|-----|----|----|----|----|-------------|-------------|
| LM  | *  | 12 | 3  | *  | *           | 100%        |
| LAD | 10 | 1  | 1  | 3  | 90.9%       | 25.0%       |
| LCX | 4  | 7  | 2  | 2  | 66.7%       | 77.8%       |
| RCA | 2  | 8  | 2  | 3  | 50.0%       | 72.7%       |

\*There was no positive results reported in ICA for LM.





Fig 5: ROC curve of CCTA vs ICA in  $\geq 50\%$  (A), and  $\geq 70\%$  (B) stenosis. Both thresholds reveal the same area under curve of 0.5333 with a confidence interval of 0.3238 to 0.7429

### Finding 2: CTCS vs ICA

Fig 6: ROC for identifying significant ( $\geq 50\%$ ) stenosis by CTCS with  $\geq 400$  threshold showing high detection accuracy of significant stenosis with urea under curve of 0.9615.



## CONCLUSIONS

Our results indicate that CTCS is a reliable technique to detect significant coronary stenosis as agreeing with [1] finding. Hence it has the potential to be used as a predicting method for stenosis in symptomatic patients before invasive procedures. Demonstrated also is a significant stenosis measurements in CTCS which correlates better than CCTA with ICA. Additionally, there was weak correlation between CCTA and ICA. However, CCTA detection accuracy might be better in newly formed coronary lesions than the detection of significant chronic coronary stenosis. This observation reasonably agrees with [4 and 5].

## ACKNOWLEDGMENT

We would like to gratefully acknowledge the staff in the Department of Medical Diagnostic Imaging at the University Hospital Sharjah. We would like to express our deepest appreciation to Dr. Abdulmunheem Obaideen (Consultant Radiologist and Chair), Dr. Sammy Weis (Senior Specialist Radiologist), Omar Adas (Radiation Protection Officer and Senior Radiology Technologist), and Fares Abdulghani (Radiology Technologist) for their help and support that made this study accomplishable.

## SELECTED REFERENCES

- [1] Lau et al. (2005). Radiology, 235(2), 415-422.
- [2] Noncommunicable Diseases Country Profiles WHO. (2014) (1st ed.). Switzerland
- [3] Schoenhagen et al (2004). Radiology, 232(1), pp. 7–17.
- [4] Nieman et al. (2001). The Lancet, 357(9256), 599-603.
- [5] Krupiński et al. (2012). Clinical Imaging, 36(6), 724-731.